

Expanded BRFSS, 2003

Richmond Locality

Table of Contents:

Optional Module 1: Adult Asthma History 1
Optional Module 5: Diabetes..... 2
Optional Module 7: Disability/Quality of Life..... 6

Optional Module 10: Health Care Coverage 10
Optional Module 13: Injury Control - Falls..... 12
Optional Module 19: Tobacco Consumption 13
Optional Module 20: Tobacco - Media 14
Optional Module 21: Tobacco – other products 22
Optional Module 22: Tobacco – Work site ETS 23
Optional Module 23: Seat Belts..... 25
Optional Module 24: Weight Control..... 27

New York State Department of Health

Expanded BRFSS, 2003 - New York State Department of Health

Have had an Asthma Episode or Attack in the Past 12 Months, among Those with Asthma

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	32	61.8	18	38.2	15.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Check Blood for Glucose or Sugar at Least Once per Day among Those with Diabetes

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	26	69.0	15	31.0	15.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Check Feet for Sores or Irritations at Least Once per Day among Those with Diabetes

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	25	61.6	13	38.4	17.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had Feet Checked by a Medical Professional in Past Twelve Months, among Those with Diabetes

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	27	65.6	12	34.4	17.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had Dilated Eye Exam in Past 12 Months, among Those with Diabetes

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	30	74.5	11	25.5	14.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with a Disability

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	96	13.5	518	86.5	2.9
Gender					
Male	40	13.8	214	86.2	4.5
Female	56	13.2	304	86.8	3.7
Age					
18-34	12	6.7	128	93.3	4.1
35-54	43	15.6	221	84.4	4.8
≥55	40	18.1	157	81.9	5.8
Education					
≤High School	47	17.2	198	82.8	5.2
>High School	48	11.2	318	88.8	3.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with a Disability Requiring Help

Richmond

Demographic Groups	n ¹	Disability need help % ²	C.I. ³	n	Disability no help needed %	C.I.	n	No Disability %	C.I.
Total	43	6.2	2.0	53	7.3	2.1	518	86.5	2.9
Gender									
Male	16	5.7	3.0	24	8.1	3.5	214	86.2	4.5
Female	27	6.7	2.7	29	6.6	2.6	304	86.8	3.7
Age									
18-64	39	6.8	2.3	39	6.4	2.2	418	86.8	3.1
≥65	4	3.7	4.1	13	11.7	6.6	88	84.6	7.5
Education									
≤High School	25	9.6	4.1	22	7.6	3.5	198	82.8	5.2
>High School	18	4.2	2.1	30	7.0	2.7	318	88.8	3.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

This page left intentionally blank

This page left intentionally blank

Expanded BRFSS, 2003 - New York State Department of Health

Main Reason for Not Having Health Care Coverage

Richmond

Demographic Groups	n ¹	Changed or lost job % ²	C.I. ³	n	Could not afford %	C.I.	n	Other %	C.I.
Total	5	9.4	8.3	11	22.3	15.7	26	68.2	16.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

No Health Care Coverage for Past Two Years Among Uninsured

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	19	42.4	25	57.6	17.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Fall in Past Three Months

Richmond

Demographic Groups	n ¹	No fall % ²	C.I. ³	n	Fall with injury %	C.I.	n	Fall no injury %	C.I.
Total	271	84.6	4.8	11	3.8	2.6	36	11.6	4.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Number of Cigarettes Smoked per day among Everyday Smokers

Richmond

Demographic Groups	n¹	1-10 cigarettes %²	n	≥11 cigarettes %	C.I.³
Total	38	37.5	63	62.5	11.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Television

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	361	62.2	221	37.8	4.6
Gender					
Male	146	59.9	94	40.1	7.3
Female	215	64.3	127	35.7	5.7
Age					
18-34	91	64.4	44	35.6	9.4
35-54	153	60.3	99	39.7	6.8
≥55	112	62.8	74	37.2	8.0
Education					
≤High School	150	63.8	89	36.2	7.1
>High School	210	61.5	130	38.5	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Radio

Richmond

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	200	36.0	383	64.0	4.5
Gender					
Male	95	40.7	147	59.3	7.2
Female	105	31.8	236	68.2	5.6
Age					
18-34	57	41.5	78	58.5	9.4
35-54	90	37.2	162	62.8	6.8
≥55	51	28.8	136	71.2	7.6
Education					
≤High School	75	34.5	160	65.5	7.2
>High School	124	37.1	221	62.9	5.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Billboards

Richmond

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	218	37.1	365	62.9	4.5
Gender					
Male	92	39.6	150	60.4	7.2
Female	126	34.8	215	65.2	5.7
Age					
18-34	68	49.9	66	50.1	9.6
35-54	96	34.5	156	65.5	6.5
≥55	49	26.3	138	73.7	7.3
Education					
≤High School	81	33.0	154	67.0	7.0
>High School	137	39.9	208	60.1	5.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Buses

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	147	25.9	434	74.1	4.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Magazines

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	304	49.8	291	50.2	4.6
Gender					
Male	135	52.7	112	47.3	7.2
Female	169	47.3	179	52.7	5.9
Age					
18-34	61	44.6	74	55.4	9.5
35-54	134	48.9	124	51.1	6.8
≥55	105	57.5	86	42.5	8.0
Education					
≤High School	127	50.7	111	49.3	7.4
>High School	176	49.5	178	50.5	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from a Website

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	46	8.3	550	91.7	2.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Movie Theater

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	63	10.7	536	89.3	3.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard or Seen Anti-Tobacco Advertisement that Refers to the NYS Quitline in Past 30 Days

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	220	39.7	349	60.3	4.6
Gender					
Male	100	41.7	140	58.3	7.2
Female	120	37.9	209	62.1	6.0
Age					
18-34	67	48.4	66	51.6	9.6
35-54	96	39.4	148	60.6	6.8
≥55	55	31.5	126	68.5	8.0
Education					
≤High School	80	36.9	146	63.1	7.4
>High School	140	41.7	200	58.3	6.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Used or Tried Any Smokeless Tobacco Products such as Chewing Tobacco or Snuff

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	34	7.3	567	92.7	2.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Work Indoors Most of the Time

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	278	76.3	76	23.7	5.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Smoking Policy at Place of Work among Indoor Workers*Richmond*

Demographic Groups	n¹	Not allowed anywhere %²	C.I.³	n	Allowed in some or all places %	C.I.	n	No official policy %	C.I.
Total	248	88.0	5.1	21	9.1	4.3	6	2.8	3.0
Gender									
Male	94	84.9	9.1	9	10.1	7.3	4	4.9	6.2
Female	154	90.7	5.0	12	8.3	4.8	2	1.0	1.6
Age									
18-34	61	86.1	10.4	5	7.5	6.7	3	6.4	8.7
35-54	141	91.5	5.0	9	7.0	4.7	3	1.5	1.8
≥55	41	83.0	16.1	6	17.0	16.1	0	0.0	0.0
Education									
≤High School	74	81.5	12.2	6	10.1	8.2	4	8.4	10.4
>High School	173	90.3	5.1	15	8.8	5.0	2	0.8	1.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Use Seatbelt when Ride or Drive in a Car (Always, Nearly Always or Never Ride in a Car)

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	535	88.1	65	11.9	3.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Use Seatbelt when Ride or Drive in a Car (Always or Never ride in a Car)

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	452	74.5	148	25.5	4.1
Gender					
Male	180	69.8	67	30.2	6.9
Female	272	78.7	81	21.3	4.6
Age					
18-64	363	73.9	122	26.1	4.6
≥65	82	79.8	22	20.2	8.7
Education					
≤High School	169	70.2	72	29.8	6.8
>High School	280	77.1	76	22.9	5.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Trying to Lose Weight

Richmond

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	312	50.9	286	49.1	4.6
Gender					
Male	100	41.0	145	59.0	7.1
Female	212	59.6	141	40.4	5.8
Age					
18-34	58	39.8	78	60.2	9.2
35-54	154	59.7	104	40.3	6.7
≥55	94	49.8	99	50.2	8.2
Education					
≤High School	117	46.9	124	53.1	7.3
>High School	193	53.3	161	46.7	5.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Trying to Maintain Current Weight of Those not Trying to Lose Weight

Richmond

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	172	56.7	114	43.3	6.7
Gender					
Male	83	53.7	63	46.3	9.4
Female	89	60.6	51	39.4	9.4
Age					
18-34	36	41.8	42	58.2	12.2
35-54	73	74.0	31	26.0	9.0
≥55	59	55.9	40	44.1	11.2
Education					
≤High School	67	44.6	57	55.4	10.1
>High School	105	65.5	56	34.5	8.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Eating Fewer Calories or Less Fat to Lose or Maintain Weight

Richmond

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	377	79.3	100	20.7	4.1
Gender					
Male	126	73.2	54	26.8	7.2
Female	251	83.9	46	16.1	4.8
Age					
18-34	68	71.4	24	28.6	10.4
35-54	195	84.7	31	15.3	5.5
≥55	105	76.7	44	23.3	7.0
Education					
≤High School	143	80.0	41	20.0	6.3
>High School	233	79.2	58	20.8	5.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Using Physical Exercise to Lose or Maintain Weight

Richmond

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	321	68.5	160	31.5	4.7
Gender					
Male	123	70.1	60	29.9	7.3
Female	198	67.2	100	32.8	6.0
Age					
18-34	74	76.7	19	23.3	9.6
35-54	153	69.5	73	30.5	6.6
≥55	87	59.9	65	40.1	8.9
Education					
≤High School	106	59.7	77	40.3	8.1
>High School	215	73.5	81	26.5	5.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval